COLONOSCOPY Date:			Procedure location: 3A-166A	
If you have questions or wish to cancel the procedure, call 202-745-8523 Call 202-745-8522, option 5 for a recorded message of these instructions Washington DC Veterans Affairs Medical Center 50 Irving St., NW, Washington, DC 20422				
7 DAYS BEFORE TEST	3 & 2 DAYS		5.4V.6T.TEG	
	BEFORE TEST	1 DAY BEFORE TEST	DAY OF TEST	
Date	Date	Date	Date	
Verify your bowel prep	Follow a LOW	Start a Clear Liquid Diet ALL DA	Y Finish your bowel prep	
(GOLYTELY)	FIBER DIET for 2	Do not eat anything solid. IF you eat a	any Start the last of your prep 6 hours before you	
If you have not received your	days	solid food, your procedure may be	leave home, you MUST finish the prep within	
bowel prep, call 202-745-8523	<u>,.</u>	rescheduled.	two hours.	
IF YOU RECEIVED Miralax (polyethylene glycol) as part of your bowel preparation,	Do not eat fruits, nuts, seeds, vegetables or red meat.	Do NOT DRINK anything that you cannot see-through or liquids that are red or purple.	*If you received 2 bottles with your prep, only drink ½ of the second bottle (OR as directed).	
START TAKING IT TWICE A DAY	Feel free to eat		YOU MUST FINISH THE PREP 4 HOURS	
	Chicken, seafood,	Prepare the bowel prep:	BEFORE YOUR PROCEDURE	
Verify your driver or escort	eggs, rice, mashed	Fill the whole Golytely bottle with	Medications:	
Your escort must check in with	potatoes, white bread	water.		
you and stay until you leave on	and pasta.	You may add Crystal Light Lemonade	Diabetics: DO NOT take your diabetes	
the test day. *COVID-19	ana pasta.	flavored powder to the Golytely.	medications. Check your blood sugar.	
pandemic changes: your escort	Drink plenty of clear	Refrigerate until ready to use.		
must pick you up within 30	liquids		DO NOT take diuretics (WATER PILLS).	
minutes of your discharge time.		Drink the bowel prep at 6 pm:		
		If you have 1 bottle of the GOLYTELY		
BLOOD-THINNING		drink HALF of the bottle.	medications with small sips of water such as:	
MEDICATIONS: May need to be		If you have 2 bottles, drink 1 entire	Blood pressure, anti-arrhythmia (heart), pain	
stopped. You must contact your		bottle.	medication.	
Primary care Physician if you take		Drink 1 glass every 10 minutes until		
any of these medications:		finished.	REPORT TO GI CHECK-IN, ROOM 3A-166A	
Warfarin [Coumadin]		Continue to drink plenty of water unt	1 hour prior to your appointment time	
Clopidogrel [Plavix]		midnight.		
Rivaroxaban [Xarelto]			An adult must be present with you at check-in	
Apixaban [Eliquis]		Medications:	and stay until you leave. Your procedure will	

Diabetics: take half dose of diabetic

twice daily.

medication in the evening **IF** you take it

be rescheduled if you do not have an adult with

you. *COVID-19 pandemic changes: your escort

must pick you up within 30 minutes of your

discharge time.

Dabigatran [Pradaxa]

Fiber products

Dypiramidole/ASA [Aggrenox]

Stop taking the following:

Iron, Multivitamins, Metamucil or